

Poultry FeedChar

pastus⁺



Natural materials: wood, organic spelt husks, fruit stones

Effects: The beneficial effects of supplements made from vegetable carbons in animal diet have been known for many years. The large surface and the associated physical effects of the vegetable carbon have a positive impact on the digestive tract and thus indirectly on the animal's general well-being.

Dosage: 0.5-1.0% of the feed weight per day (or e.g.: 5 chicken = 1 measuring spoon/day).

Composition: FeedChar is produced by a mixture of different plant carbons which are optimized for your animal.

Feed material – store dry!

www.char-line.com

Heavy metals:

arsenic (As):	< 0,8 mg/kg
cadmium (Cd):	< 0,2 mg/kg
lead (Pb):	< 2 mg/kg
mercury (HG):	< 0,07 mg/kg

PAH (2)

- Controlled by the AGES (1) in accordance with the feed regulation
- PAHs are more likely to be bound through CharLine FeedChar than released".

Dioxins

Tested in accordance with Regulation (EU) Nr. 744/2012 as amended.

All vegetable carbons used in a certain mixing ratio for the FeedChars of the CharLine company have been tested.

According to the feed regulation and the underlying testing method, there is no measurable amount of PAH. The original lab report is available for download on our website.

Limit: 0.75 ng/kg TEQ.

< 0,3 ng/kg TEQ

Sum of dioxins and dioxin-like PCBs

(1) AGES = Austrian Agency for Health and Food Safety
(2) PAH = Polycyclic Aromatic Hydrocarbons

crude fibre:	0 %
crude fats:	< 0,4 %
crude ash:	15,4 %
moisture:	30 %
calcium:	0,7 %
phosphor	0,17 %

Use of medication

You should suspend the application of FeedChar during the use of medication, as there is a possibility of binding active substances.

version: 2023_01